

Learning Goal: I will be able to use the properties of the Cartesian equation to solve problems and convert between different equations of planes.

Minds On: Different Types of Eq`ns

Action: Note and practice

Consolidation: Exit Question

Minds On

Picture it!

Using a 3D model (such as your textbook) and a plane (such as this piece of paper), you can picture a line L that is perpendicular to the plane, π . For any plane in \mathbb{R}^3 , there is only one possible line that can be drawn through the origin perpendicular to the plane. This line is called the **normal axis** for the plane, and the direction vector is called the **normal** to the plane. The direction of the normal axis is given by a vector joining the origin to any point on the normal axis. This means there are an infinite number of normals for any plane.

Minds On

Cartesian Equation of a Plane

The Cartesian (or scalar) equation of a plane in \mathbb{R}^3 is of the form $Ax + By + Cz + D = 0$, with normal $\vec{n} = (A, B, C)$. The normal \vec{n} is a nonzero vector perpendicular to all vectors in the plane.

Minds On

Try on your own or with a partner: The point $A(1, 2, 2)$ is a point on the plane with normal $\vec{n} = (-1, 2, 6)$. Determine the Cartesian equation of this plane.

$$Ax + By + Cz + D = 0$$

$$(-1)x + (2)y + (6)z + D = 0$$

$$(-1)(1) + (2)(2) + (6)(2) + D = 0$$

$$-1 + 4 + 12 + D = 0$$

$$D = -15$$

$$\therefore -x + 2y + 6z - 15 = 0$$

or

$$x - 2y - 6z + 15 = 0$$

A should be \oplus

Action

Example 1: Determine the Cartesian equation of the plane containing the points $A(-1, 2, 5)$, $B(3, 2, 4)$, and $C(-2, -3, 6)$.

We need normal + point.

Find direction vectors \vec{m}_1 and \vec{m}_2

$$\begin{aligned}\vec{m}_1 = \vec{AB} &= (3 - (-1), 2 - (2), 4 - (5)) \\ &= (4, 0, -1)\end{aligned}$$

$$\vec{m}_2 = \vec{AC} = (-1, -5, 1)$$

Now.... find $\vec{n} = \vec{m}_1 \times \vec{m}_2$

next page

$$\begin{array}{r}
 \begin{array}{ccc}
 0 & \nearrow & -5 \\
 -1 & \searrow & 1 \\
 4 & \nearrow & -1 \\
 0 & \searrow & -5
 \end{array} &
 \begin{array}{l}
 x = (0) - (-5) \\
 = -5 \\
 y = (1) - (4) \\
 = -3 \\
 z = (-20) - (0) \\
 = -20
 \end{array}
 \end{array}$$

$$\vec{n} = (-5, -3, -20)$$

$$\vec{n} = -1(5, 3, 20)$$

our normal will be $(5, 3, 20)$

$$Ax + By + Cz + D = 0$$

$$5x + 3y + 20z + D = 0$$

Find D

Sub in a given point
I'll use B (Ally said all ⊕)

$$5(3) + 3(2) + 20(4) + D = 0$$

$$15 + 6 + 80 + D = 0$$

$$D = -101$$

$$\therefore 5x + 3y + 20z - 101 = 0$$

Action

Example 2: Determine the vector and parametric equations of the plane with Cartesian equation $x - 2y + 5z - 6 = 0$. Perform a check to verify your answer.

Substitute $y=s$ and $z=t$ $\vec{n} = (1, -2, 5)$

$$x - 2s + 5t - 6 = 0$$

$$x = 2s - 5t + 6$$

parametric

$$\begin{cases} x = 6 + 2s - 5t \\ y = s \\ z = t \end{cases}$$

vector

$$\begin{cases} y = 0 + 1s + 0t \\ z = 0 + 0s + 1t \end{cases}$$

$$\vec{r} = (6, 0, 0) + s(2, 1, 0) + t(-5, 0, 1)$$

not sufficient evidence
If $\vec{n} \perp \vec{m}_1$ and
 $\vec{n} \perp \vec{m}_2$ our
normal is \perp to both direction
vectors

$$\begin{aligned}\vec{n} \cdot \vec{m}_1 &= (1, -2, 5) \cdot (2, 1, 0) \\ &= (2) + (-2) + (0) \\ &= 0 \checkmark\end{aligned}$$

$$\begin{aligned}\vec{n} \cdot \vec{m}_2 &= (1, -2, 5) \cdot (-5, 0, 1) \\ &= (-5) + (0) + (5) \\ &= 0 \checkmark\end{aligned}$$

find $\vec{m}_1 \times \vec{m}_2$
 (will give us a normal)

$$\begin{array}{l} 1 \quad \swarrow \searrow \quad 0 \quad x=1 \\ 0 \quad \swarrow \searrow \quad 1 \\ 2 \quad \swarrow \searrow \quad -5 \quad y=-2 \\ 1 \quad \swarrow \searrow \quad 0 \quad z=5 \end{array}$$

$$\vec{m}_1 \times \vec{m}_2 = (1, -2, 5)$$

that's our normal!

* This is equivalent to
 the dot products we did.

Do this OR the dot products
 on previous page.
 THEN, verify D

Verify D

$$x - 2y + 5z - 6 = 0 \quad (\text{given})$$

$$\vec{r} = (6, 0, 0) + s(2, 1, 0) + t(-5, 0, 1)$$

(found)

Test \vec{r}_0 in given equation
to show point is on plane,

$$\begin{aligned} \text{LS} & & \text{RS} \\ (6) - 2(0) + 5(0) - 6 & \stackrel{?}{=} 0 \\ & = 6 - 0 + 0 - 6 \\ & = 6 - 6 \\ & = 0 \end{aligned}$$

equations are equivalent

Action

Parallel and Perpendicular Planes – Picture it!

1. If π_1 and π_2 are two perpendicular planes, with normals \vec{n}_1 and \vec{n}_2 , respectively, their normals are perpendicular (that is, the dot product is zero).
 2. If π_1 and π_2 are two parallel planes, with normals \vec{n}_1 and \vec{n}_2 , respectively, their normals are parallel (that is $\vec{n}_1 = k\vec{n}_2$) for all nonzero real numbers k .
-

Action

Angle between Intersecting Planes

The angle, θ , between two planes, π_1 and π_2 , with normals of \vec{n}_1 and \vec{n}_2 , respectively, can be calculated using the formula $\cos\theta = \frac{\vec{n}_1 \cdot \vec{n}_2}{|\vec{n}_1||\vec{n}_2|}$.

Action

Example 3: Determine the acute and obtuse angle between the two planes

$\pi_1: x - y - 2z + 3 = 0$ and $\pi_2: 2x + y - z + 2 = 0$.

$$\vec{n}_1 = (1, -1, -2) \quad \vec{n}_2 = (2, 1, -1)$$

$$\cos \theta = \frac{\vec{n}_1 \cdot \vec{n}_2}{|\vec{n}_1| |\vec{n}_2|}$$

$$\cos \theta = \frac{(1, -1, -2) \cdot (2, 1, -1)}{(\sqrt{1^2 + (-1)^2 + (-2)^2})(\sqrt{2^2 + 1^2 + (-1)^2})}$$

$$\cos \theta = \frac{(2) + (-1) + (2)}{(\sqrt{6})(\sqrt{6})}$$

$$\cos \theta = \frac{3}{6}$$

$$\cos \theta = \frac{1}{2}$$

$\therefore \theta = 60^\circ$
and obtuse angle = 120°

Consolidation

- a) Show that the planes $\pi_1: 2x - 3y + z - 1 = 0$ and $\pi_2: 4x - 3y - 17z = 0$ are perpendicular.
- b) Show that the planes $\pi_3: 2x - 3y + 2z - 1 = 0$ and $\pi_4: 2x - 3y + 2z - 3 = 0$ are parallel but not coincide

a) $\vec{n}_1 = (2, -3, 1)$
 $\vec{n}_2 = (4, -3, -17)$
 if \perp , $\vec{n}_1 \cdot \vec{n}_2 = 0$

$$\begin{aligned} \vec{n}_1 \cdot \vec{n}_2 &= (2, -3, 1) \cdot (4, -3, -17) \\ &= (8) + (9) + (-17) \\ &= 17 - 17 \\ &= 0 \checkmark \end{aligned}$$

$$\therefore \pi_1 \perp \pi_2$$

$$\begin{aligned} \text{b) } \vec{n}_1 &= (2, -3, 2) & \vec{n}_1 &= \vec{n}_2 \\ \vec{n}_2 &= (2, -3, 2) \end{aligned}$$

$$\therefore \Pi_3 \nparallel \Pi_4$$

$$\text{but } D_1 \neq D_2$$

$$\therefore \text{non-coincident}$$

Need to know

Equation Table

• product $\rightarrow \cos\theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$

x product \rightarrow how to

if $\vec{a} \cdot \vec{b} = 0$, then $\vec{a} \perp \vec{b}$

$\vec{a} \times \vec{b} = \vec{c} \rightarrow \vec{a} \perp \vec{c}$ AND $\vec{b} \perp \vec{c}$

Requirements for a plane